

# Chapter 6

## DEPARTMENT OF SPACE PHYSICS

### 6.1 STAFF

#### 6.1.1 Scientific staff

Anna Antalová, Ján Baláž, Ladislav Just,  
Karel Kudela, Marián Slivka

#### 6.1.2 PhD students

Pavol Bobik (PF UPJS), Radoslav Bučík (TU  
Zvolen). Their PhD theses are lead by DSP.

#### 6.1.3 Technical staff

Vladimír Kollár, Igor Strhársky, Ronald Langer,  
Jana Štetiarová, Samuel Štefánik, Viera Štrbinová,  
Anna Tomičová

### 6.2 SCIENTIFIC ACTIVITY

#### 6.2.1 Introduction

Department of Space Physics (DSP) is one of the oldest departments of the Institute. The research is oriented to the experimental study of energetic particles in space based on measurements from the satellites (1) as well as of cosmic rays measured with the ground based instruments (2). Both types of studies are devoted mainly to the obtaining relevant informations on the physical processes within the magnetosphere of the Earth in the heliosphere. In addition, heavy nuclei interactions are studied too.

#### 6.2.2 Projects

##### Project of Slovak Scientific Agency VEGA

**G-1353** Sounding of the magnetosphere by energetic particles (1995-1997)

**G-5137** Energetic Particles in Space: Dynamics of Magnetospheric and Heliospheric Fluxes (starting from 1998)

Principal investigator: K. Kudela

The staff is participating also in other projects of VEGA lead by other institutions.

### 6.3 RESULTS

#### 6.3.1 Cosmic Ray Dynamics at Neutron Monitor Energies and above.

Neutron monitor at Lomnický štít is providing continual measurements of secondary nucleon component. The data collection, barometric pressure correction, archiving, continuous "in situ" checking of the measurements and upgrading the electronics is routinely done by the group of technicians in High Tatras (V.Kollár, R. Langer, S.Štefánik and recently also by I. Strhársky). The data are produced with 1 minute resolution. High count rate, related to the unique measurement place (2634 m altitude) is allowing to perform studies of the cosmic ray variability of small amplitudes. The review graphs of 1 month measurements together with the geomagnetic activity indices and interplanetary magnetic field are prepared by P.Bobik. The studies of solar ground level events, as well as long term cosmic ray investigations, are based on existing measurements. Recently a new registration electronic system for neutron supermonitor located on Lomnický štít observatory have been designed by I. Strhársky and colleagues. Finalization is in progress.

Cosmic ray data, obtained from numerous stations on the ground of the Earth and from underground measurements (higher energies), including in particular also those of neutron monitor at Lomnický štít, provide a large data set for examining the properties

of cosmic ray time series at various primary rigidities and covering periods corresponding to different solar activity levels. The rapporteur talk [1] summarized the recent progress and present knowledge of the selected items of quasiperiodic variations of cosmic rays and of energetic particle research relevant to the studies of terrestrial environment.

In addition to the fractal dimension estimates of cosmic ray time series we described in the preceding period, the correlation dimension of the time series of primaries about 145 GV rigidity was obtained jointly with the group of Shinshu University Cosmic Ray Group recently [2]. The influence of the diurnal variation [4] and high level white noise to the signal [3] was checked, in the same collaboration. The first experience with the description of cosmic ray time series based on wavelet transform, was obtained. The application of the method on the long term, stable measurement (Misato, Japan, underground muon telescope, 1975-1994) has shown the different behaviour of the small scale temporal variability (2 hours) and larger one ( $> 16$  hours) if compared with the solar activity level.

The relevance of cosmic ray time series for eventual prediction of the nonstationary physical processes with impact of near Earth surrounding has been checked with using Lomnický stit data [5,6]. Statistical type of studies performed here is accomplishing the single effect detailed analysis illustrating the increase of variability on a single neutron monitor before the disturbance as a consequence of increasing the cosmic ray anisotropy (Bieber and Evenson, *GRL*, 25, p.2955-2958, 1998). Thus it is reasonable to check the possibility of using the cosmic ray time series as one of the parameters of ANN (artificial neural network) and/or FNN (fuzzy neural network) for predictions of geomagnetic storms. Such approaches were used in the collaboration with PF UPJS recently on the data sets describing the plasma and magnetic field structures near the Earth [7,8,9].

It was found that the short term periodic variations of secondary cosmic rays in the Brazil magnetic anomaly region, observed by a series of balloons in that region, may be caused by the pulsating character of "screening" of primaries by the magnetosphere [10]. In addition to the periodic variation of the altitude of mirror points of trapped radiation suggested earlier as an explanation for the observed secondary variations, the variability of transparency of magnetosphere provides another possibility. Indeed the periodicities are observed also in Lomnický stit NM measurements.

The structure of forbidden and allowed rigidities for Lomnický stit and Rome stations has been

computed for various levels of geomagnetic activity described by the Kp index and using the combined IGRF (internal) and Tsyganenko 89 (external) model of the field. While the seasonal cosmic ray variation is not significant at these positions, the amplitude of the diurnal variation for the quiet time and most disturbed one ( $Kp > 6$ ) is differing by few tenths up to one per cent, assuming also the response (yield) function and the effect of the atmosphere [11,12].

Soft X-ray parameters, derived from Solar Radiation satellite measurements, were used to analyse, on daily basis, their relationship with galactic cosmic ray (GCR) modulation and geomagnetic storms. Coronal Mass Ejections (CMEs) are a principal heliospherical phenomenon, which implies a cause-effect relationship between the interplanetary shocks responsible for GCR-modulation as well as geomagnetic storms. The main difficulty in investigating the solar origin of the nonrecurrent heliospherical shocks arises from the fact that the plasma clouds ejected by the Sun are, in general, observable with the coronagraphs only when they occur near the limb of the Sun. Thus, the detection of CMEs on the disk must rely on some proxy parameter. One of the most convenient appears to be the intensity of the solar soft X-ray flare emission, and in particular those called Long Duration Events (LDEs). The analysis of soft X-ray data yields the following results: We investigate [13] the role of the coronal mass ejections (by using soft X-ray solar emission as proxy data) in producing the nonrecurrent geomagnetic storms in the period 1969-1974. The linkage between these phenomena is confirmed however, in turns out that CMEs associated with chromospheric flares, accompanied by type IV radio emission, are the most effective in perturbing the geomagnetic field. From the multivariate daily sequence analyses of GCR/Solar X-Ray Parameters [14] it was found that extreme values of the cross-correlation functions (on daily basis) remain less than 0.5. The periods in which these values are significant correspond to the ones in which large-scale activity phenomena occur on the Sun (in 1970.5-1971.5 and 1972 years). We examine the short-term variability of GCR records, from two neutron monitors with different cut-off rigidities, namely Calgary (CA) and Lomnický stit in 1968–1996 years. As for the LDE flare index, the new consecutive daily sequences were calculated on the daily basis [15-18]. The autocorrelation analysis [19] of the SXR solar parameters (XBG and LDE) clearly shows, that the behavior of the medium-term X-corona (i.e. within a 27-day synodic solar rotation) depends on the solar cycle phase. Instead, on a short-term scale (1 day) this dependence exists

only for the X-flaring corona, with the X-background nearly stable from the maximum to the minimum activity phases.

collected by **K. Kudela and A. Antalová**

### 6.3.2 Medium Energy particles within the Magnetosphere and near its Boundaries.

Populations of particles with the energies well below that of cosmic rays and well above the solar wind (from few tens of keV up to several MeV) are systematically studied by the DSP in last few years. The analysis of the obtained data from both low altitude and high apogee satellites, as well as development and construction of new instruments for the future studies have been carried out during last two years.

The Interball project was realized by launch of two satellites (Interball-1, Interball-2) and two corresponding subsatellites (Magion 4 and Magion 5) in 1995 and 1996, respectively. The measurements with DOK2/DOKS instruments provided large amount of data on energy spectra and angular distribution of energetic particles in the outer magnetosphere, in the geomagnetic tail, within the magnetosheath and in the region upstream of the bow shock. The amount of the data (for Interball-1 now more than three years partly covered by DOK2, [33,34]) required large effort in data processing and archiving (done especially by M.Slivka and J. Štetiarová). The data processing contains the separation of the frame modes (time profiles and energy spectra of electrons and protons), and merging the particle flux data with the time, satellite position, IMF vector, geometry of connection to the bow shock (in the region upstream of the bow shock), orientation and pitch angles of single detectors. These works are in progress. For reviewing the data as well as for the multi-instrument and multi-satellite studies the review pictures are constructed. The comparison of two point measurements of energetic particles in wide energy range (Interball 1 and Magion 4) have confirmed the result found by us earlier on the data from Prognoz-10 [26]. The statistical studies of the ion and electron flux within the magnetosheath based on Prognoz-10 and Interball 1 measurements have indicated that energy spectra of the particles close to the magnetopause are harder than those deep in that region [43]. The streaming of ions dawnward and duskward close to the magnetopause is not showing any clear pattern in (ZY) GSE plane. This is consistent with suggestion of patchy structure of merging at the magnetopause and with the leakage

of magnetospheric particles to magnetosheath. The fluxes of energetic ions at Interball 1 upstream from the bow shock for the positions of the satellite not connected to the bow shock along the field line (simplified assumption of the field line as a straight line) are providing the upper limit to the quiet time fluxes in the heliosphere studied in detail [27]. Several particular results have been presented at the NATO Advances Research Workshop "Coordinated Studies of the Solar Wind-Magnetosphere-Ionosphere Interaction: Interball observations" held in Kosice, September 7 - 11, 1998. The abstracts of 7 papers with our coauthorship are in the abstract booklet.

Low altitude measurements with use of SPE1 instrument onboard the Active satellite have shown strong, short duration (few tenths of second) pulsations of electrons precipitating to the atmosphere. The patterns of these pulsations in subauroral and auroral regions have been described in [22,24,25]. Both pitch angle and energy diffusion is deduced from them. The pulsations are found in the regions with enhanced intensity of VLF emissions having also spiky character.

For the low altitude satellite COMPASS (checking the eventual precursors of the earthquake activity in magnetospheric data at low altitudes) the monitor of energetic particles MEP-1 [31] using the silicon detectors and highly flexible system of data recording has been developed in the cooperation with IEP RAS Moscow and UNICAMP Campinas. The technical group of the DSP developed and finalized programmable particle spectrometer MEP-1 [28,29] for this satellite. The flight model of the MEP-1 is ready for the launch.

Some theoretical and practical methods and tools were developed for system identification of particle registration devices (geometrical factor, 3D-anisotropy) and for study of system dynamic behaviour by programmable stochastic photostimulation [30].

The distribution of gamma ray flux based on the whole set of available data from CORONAS-I (experiment SONG) was obtained in geographic coordinates showing the strong increase in the brazil magnetic anomaly region [32].

collected by **K. Kudela, M. Slivka and J. Baláz**

### 6.3.3 Fragmentation of Relativistic Nuclei in Nuclear Emulsion

The knowledge of fragmentation parameters in nuclear collisions is important in particular for the understanding of the composition of cosmic rays in its

sources, for the transport as well as for studies of galactic cosmic ray exposure behind the shielding in space flights. Recently the interactions of  $^{84}\text{Kr}$  at energy 0.95 GeV/nucleon with nuclear photoemulsion have been studied. The bounce-off effect of primary nuclei fragments were confirmed using the azimuthal correlation function method. The magnitude of this effect depends on the impact parameter of interacting nuclei. This effect is independent on the magnitude of primary energy in the region up to 2 GeV/nucleon. [35,41].

The angular distribution of projectile-associated He fragment from Pb induced reactions on stationary Pb target at 158 A GeV/c has been studied. Two emission components are appearing in the projected angular spectra of the He particles [36]. The charge distribution of relativistic nuclei fragmenting (with  $Z > 2$ ) of  $^{197}\text{Au}$  nuclei at 10.7 GeV/nucleon has been measured. With the help of moments of these distributions research, we found out the asymptotic behavior of fragmentation's parameters for the peripheral collisions of  $^{197}\text{Au}$  nuclei in emulsion. [39].

With the help of transverse-momentum of relativistic fragments of primary nuclei study where registered collisions with Ag(Br) nuclei at energy 10.7 GeV/nucleon. Its correlation with the emission of target nuclei fragment were studied too [40]. We observed found out large cross section for fission Au+Emul. in comparison with published. After this analysis could calculate the fission yields of Au nuclei on different nuclear targets. The fission yield for Au+H is  $Y_H = 0.041 \pm 0.025$ , in agreement with the value obtained from the p+Au in wide range energies. For the interactions on the CNO group we obtained  $Y_{CNO} = 0.026 \pm 0.018$  and for AgBr  $Y_{AgBr} = (5.14 \pm 2.6) \cdot 10^{-3}$ . [42] The results obtained in the EMU01 collaboration include the coauthors from Armenia, Australia, China, Germany, India, Kazakhstan, Russia, Slovakia, Sweden, USA, Uzbekistan.

collected by L. Just

## 6.4 SELECTED PAPERS

1. Kudela K., Quasiperiodic Variations and Terrestrial Environment, Rapporteur paper, in vol. 8 of the 25th ICRC, Durban, S.Africa, p. 175 - 192, World Scientific Co. Pte. Ltd., Singapore-New Jersey-London-Hong Kong, ed. M.S. Potgieter, B.C. Raubenheimer, D. J. van der Walt, 1998

2. Yasue S., K. Munakata, T. Shiozawa, A.Ogi,

K. Kudela: Influences of the solar activity on the correlation dimension of underground muon time series at 34 mwe, Accepted for publication in *Advances in Space Res.*, 1998

3. Yasue S., T. Yamazaki, K. Munakata, C. Kato, D.L.Hall, S. Mori, K. Kudela: Evaluation of the valid fractal dimension of cosmic ray time series: I. Including the High Level Background White Noise, Proc. 25th ICRC, Durban, S.Africa, vol. 2, p. 213-216, 1997

4. Yasue S., K. Munakata, T. Yamazaki, A. Ogi, T. Shiozawa, K. Kudela: Evaluation of the Valid Fractal Dimension of Cosmic Ray Time Series: II. Including the Marked Periodic Variation, Proc. 25th ICRC, Durban, S. Africa, vol. 2, p. 209-212, 1997

5. Kudela K., G. Andrejkova, S. Yasue, K. Munakata: Cosmic Ray Temporal Variability: Potential Relevance for Geomagnetic Activity Predictions?, Proc. of a Workshop Solar Terrestrial Predictions V, RWC Tokyo, Hiraiso Solar Terrestrial Research Center, Communications Research Laboratory, Hitachinaka, Ibaraki, Japan, p. 244-247, November 1997

6. Kudela K., E.O. Fluckiger, R. Langer, P. Bobik, Power spectra of neutron monitor time series at frequencies  $f > 1.15 \cdot 10^{-5}$  Hz, Proc. 25th ICRC, Durban, S. Africa, vol. 2, p. 425-427, 1997

7. Andrejkova G., H. Toth, K. Kudela, Fuzzy Neural Networks in the Predictions of Geomagnetic Storms, European Space Agency WPP-148, p. 173-179, 1997

8. Andrejkova G., K. Marcisinova, K. Kudela, Genetic Algorithms in the Prediction of Geomagnetic Storms, European Space Agency WPP-148, p. 181-186, 1997

9. Andrejkova G., H. Toth, K. Kudela, The Approach of Fuzzy Neural Networks for Geomagnetic Storm Predictions, ESA Workshop on Space Weather, ESTEC, The Netherlands, November 1998 (contribution accepted for publication)

10. Martin I.M., A.A. Gusev, K. Kudela, J. Kassovicova, G.I. Pugacheva, On the Periodic Variations of Secondary Cosmic Rays and the Geomagnetic Pc4 Pulsations in Brazil Magnetic Anomaly Region, Ann. Geophysicae, Atmospheres, Hydrosphere and Space Science, vol.15, p.1008-1014, 1997

11. Kudela K., M. Storini, P. Bobik, J. Kassovicova, Access of Cosmic Rays to Lomnický štít and Rome stations, paper presented at 16th ECRS (European Cosmic Ray Symposium), Alcalá de Henares, Spain, July 20-24, 1998
12. Bobik P., K. Kudela, M. Storini, J. Kassovicova, Access of Cosmic Rays to Lomnický štít and Rome stations, Proc. of Contributed Papers, WDS 98 (Week of doctoral students 98), Fac. Math. Physics, Charles U., Prague, ed. J. Safrankova, Part II, Physics of Plasmas and Ionized Media, p. 234-239, 1998
13. Landi, R., Moreno, G., Storini, M. and Antalová, A., Coronal mass ejections, flares, and geomagnetic storms, Journal of Geophysical Research 103, A9, 20553-20559, 1998.
14. Jakimiec, M., Storini, M. and Antalová, A., Multivariate daily sequence analyses of GCR/Solar X-ray parameters, Adv. Space Res. 20, 111-114, 1997.
15. Antalová, A., 1977 and 1978 solar soft X-ray daily flare indices, Contrib. Astron. Obs. Skalnaté Pleso 28, 63-72, 1998
16. Antalová, A., The continuation of the daily soft X-ray flare index (1973-1974), Contrib. Astron. Obs. Skalnaté Pleso 27, 31-40, 1997
17. Antalová, A. The 20-th solar cycle minimum and its daily soft X-ray flare index (1975-1976), Contr. Astron. Obs. Skalnaté Pleso 27, 129-138, 1997
18. Antalová, A., 22-nd Cycle's Sectorial Distribution of LDE-type Flaring Centers, Proceedings of the Third SOLTIP, eds. X. Feng, F. Wei and M. Dryer, International Academic Publishers, Beijing 1998, 117-123., 1998
19. Jakimiec, M., Storini, M. and Antalová, A. Solar soft X-ray parameters: 1969-1976 autocorrelation functions, Contr. Astron. Obs. Skalnaté Pleso 27, 104-114, 1997
20. Storini, M., Massetti, S. and A. Antalová, To forecast huge Forbush decreases during solar activity cycles, In: 25ICRC, Contributed Papers, Vol 1, Durban, South Africa, 409-412, 1997
21. Slivka, M., K. Kudela, V.N. Lutsenko, E.T. Sarris, J. Štetiarová, NATO Advanced Research Workshop, Coordinated Studies of the Solar Wind - Magnetosphere - Ionosphere Interactions, Interball Observations, Košice, Slovakia, Sept. 7 - 11 1998, Abstract Booklet, abstract p.60, 1998.
22. Kudela, K., Slivka M., Jiricek F., Triska P., Shuiskaya F.K., Martin I.M., Strong Fluctuations of Energetic Electrons at Low Altitudes, Adv.Space Res., Vol.20, No.3, pp.499-503, 1997.
23. V.A.Sergeev, T. Bosinger, A.G. Yahnin, I.A. Kornilov, R.J. Pellinen, T.I. Pulkkinen, N.L. Borodkova, V.N. Lutsenko, M.N. Nozdrachev, V.I. Prokhorenko, A.A. Skalsky, J.A. Sauvaud, K. Kudela, M. Slivka, E.T. Sarris, Coordinated Ground Observations Near Interball-Tail Footpoint: Initial Results for Plasma Sheet Passes of Interball in 1995/1996, Proc. Third Intern. Conf. on Substorms (ICS-3), Versailles, France, 12-17 May 1996, ESA SP-386 (October 1996) pp.585-590, 1996.
24. Kudela, K., I.M. Martin, M. Slivka, F.K. Shuiskaya, Strong Pulsations of Energetic Electrons Precipitating into Atmosphere: Active Satellite Data, 25th ICRC, Durban, South Africa, Vol.2, p.341, 1997
25. Kudela, K., I.M. Martin, F. K. Shuiskaya, M.M. Mogilevsky, F. Jiricek, Pulsations of precipitating energetic electrons: Active satellite data, J. Atmos. and Solar-Terrestrial Physics, vol. 60, p. 643-653, 1998
26. Nemecek, Z., J. Safrankova, O. Santolik, K. Kudela, and E.T. Sarris, Energetic particles in the vicinity of the dawn magnetopause, Adv. Space Res., vol. 20, No 4/5, pp. 851-856, 1997
27. Kiraly, P., V. Bothmer, K. Kecskemety, E. Kepler, K. Kudela, H. Kunow, Yu. I. Ogachev, R. Mueller-Mellin, J. Rodriguez-Pacheco, R. von Steiger, J. Torsti, E. Valtonen, and M.A. Zeldovich, Search for the Origin of Quiet-time particle fluxes in the inner Heliosphere, Proc. of the 25th ICRC, Durban, South Africa, vol. 2, p. 477-480, 1997
28. Baláz, J., Strhársky, I., Programmable Particle Spectrometer MEP-1. Technical description and user's guide, 2nd edition, pp 54, IEP-SAS Košice 1998.
29. Baláz, J., Gladyshev, V., Kudela, K., Martin, I.M., Strhársky, I.: Programmable Particle Spectrometer for Space research. Accepted for publication in "Pribory i Tekhnika Eksperimenta Nr.5/98, Moscow 1998.
30. Baláz, J.: Design and system identification of energetic particle registration instruments to be flown on board a spacecraft. Doctoral thesis. pp.119. Technical University Košice 1998.

31. Kudela, K., J. Baláz, I. Strhářsky, J.Štetiarová, I.M. Martin, V. Gladyshev, F. K. Shuiskaya, A. Stepanov, V. Klepikov: MEP-1 experiment. Scientific Tasks and Basic Characteristics, Preprint IEP SAS, April 1997, 23 pp.
32. Bučík, R., K. Kudela, A.V. Bogomolov, I.N. Myagkova, S.N. Kuznetsov, S. P. Ryumin, Distribution of gamma ray fluxes at altitude 500 km: CORONAS-I data, Proc. of Week of Doctoral Students (WDS 98, Prague), ed. J.Safrankova, part II, Physics of Plasmas and Ionized Media, Fac. of Math. and Phys., Charles University, Prague, p. 216-219, 1998
33. Lutsenko, V.N., K.Kudela, E.T. Sarris, The DOK2 Experiment to Study Energetic Particles by the Tail Probe and Auroral Probe Satellites in the Interball Project, Cosmic Res., 36, No 1, p. 98 - 107, 1998
34. Taktakishvili, A.L., L.M. Zelenyi, V.N. Lutsenko, K. Kudela: On the Spectra of Energetic Particles in the Earth's Magnetotail, Cosmic Res., vol. 36, No 3, p. 282-291, 1998
35. Adamovich M.I., ..., Just L., ..., Yuldashev B.S.: Collective Motion in  $^{84}\text{Kr}$  Induced by Interactions with Ag(Br) Nuclei at 0.95 GeV per Nucleon. Jadernaja fizika, vyp.9, tom 60,1580-1585, 1997.
36. Adamovich M.I., ..., Just L., ..., Zhou D.C. (EMU01-Collab.): He production in 158 A GeV/c Pb on Pb interactions. Physics Letters B 390 , 445-449, 1997.
37. Adamovich M.I., ..., Just L., ..., Zhou D.C. (EMU01-Collab.): Charged particle multiplicities, densities and fluctuations in Pb+Pb interactions at 158 A GeV. Physics Letters B 407, 92-96, 1997.
38. Adamovich M.I., ..., Just L., ..., Zhou D.C. (EMU01-Collab.): Complex analysis of gold interactions with photoemulsion nuclei at 10.7 GeV/nucleon within the framework of cascade and FRITIOF models. Zeitschrift f. Physik A 358 , 337-351, 1997.
39. Adamovich M.I., ..., Just L., ..., Zhou D.C. (EMU 01-Collab.): Critical behaviour in Au fragmentation at 10.7 A GeV. The European Physical Journal A 1, 77-83, 1998.
40. Adamovich M.I., ..., Just L., ..., Zhou D.C. (EMU 01-Collab.): Bounce - off in  $^{197}\text{Au}$  induced collisions with Ag (Br) nuclei at 11.6 A GeV/c. The European Physical Journal A 2, 61-67, 1998.
41. Kravčáková, A., Vokál, S. and the Kr Collaboration (Dubna, Košice, Lund, Moscow, Tashkent, Varanasi): Azimuthal correlation functions as a probe for collective motion. Czechoslovak Journal of Physics, vol.48, Suppl. S1, pp. 127-132, 1998.
42. Popa V., ... Just L., ..., Zhou D.C. (EMU01-Collab.): Fission of gold nuclei at 10.6 A GeV in nuclear Emulsions. Proc. 25th ICRC, Durban, South Africa, vol.6, p. 21-24, 1997.
43. Kudela.K., D.G.Sibeck, M.Slivka, V.Lutsenko, Energetic ions in the magnetosheat: Statistical study of spectra and anisotropy. Adv.Space Res. Vol. 21, No. 4, pp. 633-636, 1998.

## 6.5 OTHER ACTIVITIES

The results described shortly in part 6.3 have been obtained in wide international collaboration which is completely listed in corresponding paragraph of the recent biennial report (domestic laboratories and those in the Czech Republic, Italy, Russia, Sweden, Switzerland, USA, Germany, Japan, Brazil, Finland, Greece and other countries).

### 6.5.1 Organization of scientific conferences

The most important activity of the international character was the NATO Advanced Research Workshop "Coordinated Studies of the Solar Wind - Magnetosphere - Ionosphere Interaction: Interball observations" organized in Košice, September 7 - 11, 1998. The director and co-director of the event were Dr. D. G. Sibeck, Johns Hopkins University, Applied Physics Laboratory, Laurel, MD, USA and K. Kudela, respectively.